

# *Ultracold quantum gases and functional renormalization*

Stefan Flörchinger (CERN)

work done in Heidelberg in collaboration with

S. Diehl, H. Gies, S. Moroz, J. M. Pawłowski, R. Schmidt,  
M. M. Scherer and C. Wetterich

Bad Honnef, 12. Februar 2011

## *Why are Ultracold quantum gases interesting?*

- Ultracold gases in the bulk are simple systems!
  - for example: Fermi surface is usually a sphere.
- Both fermions and bosons can be studied.
- Interactions can be tuned to arbitrary values.
- Lower dimensional systems can be realized.

Very nice model system to test methods of quantum and statistical *field theory*!

# Quantum field theory

- Describes also electrons, atoms, quarks, gluons, protons,...
- Crucial object: quantum effective action


$$\begin{aligned}\Gamma[\phi] &= S[\phi] + \frac{1}{2} \text{Tr} \ln S^{(2)} + \dots \\ &= \int dt \int d^d x U(\phi) + \dots\end{aligned}$$

- Quantum field equations from  $\frac{\delta\Gamma}{\delta\phi} = 0$ .
- Symmetries of  $\Gamma$  lead to conserved currents.
- All physical observables are easily obtained from  $\Gamma$ .
- $\Gamma$  is generating functional of 1-PI Feynman diagrams and depends on external parameters like  $T$ ,  $\mu$ , or  $\vec{B}$ .

## How do we obtain the quantum effective action $\Gamma[\phi]$ ?

Idea of functional renormalization:  $\Gamma[\phi] \rightarrow \Gamma_k[\phi]$

- $k$  is additional infrared cutoff parameter.
- $\Gamma_k[\phi] \rightarrow \Gamma[\phi]$  for  $k \rightarrow 0$ .
- $\Gamma_k[\phi] \rightarrow S[\phi]$  for  $k \rightarrow \infty$ .
- Dependence on  $T, \mu$  or  $\vec{B}$  trivial for  $k \rightarrow \infty$ .



## $\Gamma[\phi]$ and the grand canonical ensemble

Functional integral representation of the partition function

$$Z = e^{-\beta\Omega_G} = \text{Tr} e^{-\beta(H-\mu N)} = \int D\chi e^{-S[\chi]}.$$

Generalization with  $J = \frac{\delta}{\delta\phi}\Gamma_k[\phi]$

$$e^{-\Gamma_k[\phi]} = \int D\chi e^{-S[\phi+\chi]+J\chi-\frac{1}{2}\chi R_k \chi}.$$

- $R_k$  is an infrared cutoff function
  - suppresses all fluctuations  $R_k \rightarrow \infty$  for  $k \rightarrow \infty$ .
  - is removed  $R_k \rightarrow 0$  for  $k \rightarrow 0$ .
- $\Gamma_k[\phi]$  is the *average action* or *flowing action*.
- Grand canonical potential is obtained from  $\beta\Omega_G = \Gamma_k[\phi]$  for  $k = 0$  and  $J = 0$ .

## *How the flowing action flows*

Simple and exact flow equation (Wetterich 1993)

$$\partial_k \Gamma_k[\phi] = \frac{1}{2} \text{STr} \left( \Gamma_k^{(2)}[\phi] + R_k \right)^{-1} \partial_k R_k.$$

- Differential equation for a functional.
- For most cases not solvable exactly.
- Approximate solutions can be found from Truncations.
  - Ansatz for  $\Gamma_k$  with a finite number of parameters.
  - Derive ordinary differential equations for this parameters or couplings from the flow equation for  $\Gamma_k$ .
  - Solve these equations numerically.

# Lagrangians

We use a local field theory to describe the microscopic model.

Examples:

- 1 Bose gas with pointlike interaction

$$\mathcal{L} = \varphi^* \left( \partial_\tau - \vec{\nabla}^2 - \mu \right) \varphi + \frac{1}{2} \lambda (\varphi^* \varphi)^2.$$

- 2 Fermions in the BCS-BEC-Crossover

$$\begin{aligned} \mathcal{L} = & \psi^\dagger (\partial_\tau - \vec{\nabla}^2 - \mu) \psi + \varphi^* (\partial_\tau - \frac{1}{2} \vec{\nabla}^2 - 2\mu + \nu) \varphi \\ & - h(\varphi^* \psi_1 \psi_2 + h.c.). \end{aligned}$$

These are effective theories on the length scale of the Bohr radius or van-der-Waals length.


## *Symmetries of nonrelativistic field theories*

- U(1) for particle number conservation.
- Translations and Rotations.
- Galilean boost transformations.
- Possibly conformal symmetries.
- U(1) and Galilean invariance are broken spontaneously by a Bose-Einstein condensate.
- Galilean invariance is broken explicitly for  $T > 0$ .



## Two component Fermi gas

- Two spin (or hyperfine-spin) components  $\psi_1$  and  $\psi_2$ .
- For equal mass  $M_{\psi_1} = M_{\psi_2}$ , density  $n_{\psi_1} = n_{\psi_2}$  etc. SU(2) spin symmetry
- $s$ -wave interaction measured by scattering length  $a$ .
- Repulsive microscopic interaction: Landau Fermi liquid.
- Attractive interaction leads to many interesting effects!
- Scattering length can be tuned experimentally with Feshbach resonances.





## Truncations

For many purposes *derivative expansions* are suitable approximations. For example we use for the BCS-BEC Crossover


$$\Gamma_k = \int_{\tau, \vec{x}} \left\{ \psi^\dagger (Z_\psi \partial_\tau - Z_\psi \vec{\nabla}^2 - \mu + \Delta m_\psi) \psi \right. \\ \left. + \varphi^* (Z_\varphi \partial_\tau - A_\varphi \frac{1}{2} \vec{\nabla}^2) \varphi \right. \\ \left. - h(\varphi^* \psi_1 \psi_2 + h.c.) + \frac{1}{2} \lambda_\psi (\psi^\dagger \psi)^2 + U_k(\varphi^* \varphi, \mu) \right\}$$

- The coefficients  $Z_\varphi$ ,  $A_\varphi$ ,  $\lambda_\psi$ ,  $h$ ,  $Z_\psi$ ,  $\Delta m_\psi$  and the effective potential  $U_k$  are scale-dependent.
- The effective potential  $U_k$  contains no derivatives - describes homogeneous fields.

# Fermion self energy corrections


(Floerchinger, Scherer and Wetterich, PRA **81**, 063619 (2010))

- Flow of fermion wavefunction renormalization  $Z_\psi$



solid:  $(ak_F)^{-1} = 0$ ,  
long-dashed:  $(ak_F)^{-1} = -1$ ,  
short-dashed:  $(ak_F)^{-1} = 1$

- At the macroscopic scale  $k = 0$



## *The effective potential*

- We use a Taylor expansion around the minimum  $\rho_0$


$$U_k(\varphi^* \varphi) = -p + m^2 (\varphi^* \varphi - \rho_0) + \frac{1}{2} \lambda (\varphi^* \varphi - \rho_0)^2.$$

## The effective potential


- We use a Taylor expansion around the minimum  $\rho_0$

$$U_k(\varphi^* \varphi) = -p + m^2 (\varphi^* \varphi - \rho_0) + \frac{1}{2} \lambda (\varphi^* \varphi - \rho_0)^2.$$


- Symmetry breaking:



$$T > T_c$$



$$T < T_c$$




$$T \ll T_c$$

## The effective potential

- We use a Taylor expansion around the minimum  $\rho_0$


$$U_k(\varphi^* \varphi) = -p + m^2 (\varphi^* \varphi - \rho_0) + \frac{1}{2} \lambda (\varphi^* \varphi - \rho_0)^2.$$

- Typical flow:



## *Solving the flow equation - Phase diagram*


- Information on phase diagram is contained in form of the effective potential  $U(\rho, \mu, T)$  at macroscopic scale.
- Very nice generalization of Landau's theory!
- Example: BCS-BEC Crossover  
(Floerchinger, Scherer and Wetterich, PRA **81**, 063619 (2010).)





## Solving the flow equation - Phase diagram

- Information on phase diagram is contained in form of the effective potential  $U(\rho, \mu, T)$  at macroscopic scale.
- Very nice generalization of Landau's theory!
- Example: Superfluid Bose gas in  $d = 2$   
(Floerchinger and Wetterich, PRA 79, 013601 (2009)).




## Superfluid order in two dimensions

- Bose gas with truncation

$$\Gamma_k = \int_{\tau, \vec{x}} \left\{ \varphi^* (Z \partial_\tau - V \partial_\tau^2 - A \vec{\nabla}^2) \varphi + U(\varphi^* \varphi) \right\}$$

- Mermin-Wagner theorem: No true long range order at  $T > 0$  in  $d = 2$ .
- This implies:  $n_c = \bar{\rho}_0 \rightarrow 0$  for  $k \rightarrow 0$ .



density  $n$  (solid),  
superfluid density  $\rho_0$  (dashed),  
condensate density  $\bar{\rho}_0$  (dotted),

# *Solving the flow equation - Thermodynamic observables*

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))

# Solving the flow equation - Thermodynamic observables


From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))

- entropy density  $s = -\frac{\partial U}{\partial T}$ ,




# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,

- energy density

$$\epsilon = -p + Ts + \mu n,$$


# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,


# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,
- isoth. compressibility  $\kappa_T$ ,


# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,
- isoth. compressibility  $\kappa_T$ ,
- adiab. compressibility  $\kappa_S$ ,




# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,
- isoth. compressibility  $\kappa_T$ ,
- adiab. compressibility  $\kappa_S$ ,
- velocity of sound  $l$ ,


# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,
- isoth. compressibility  $\kappa_T$ ,
- adiab. compressibility  $\kappa_S$ ,
- velocity of sound I,
- velocity of sound II,


# Solving the flow equation - Thermodynamic observables

From grand canonical potential

$$dU = -dp = -s dT - n d\mu$$

take derivatives e. g. for Bose gas in  $d = 3$

(Floerchinger and Wetterich, PRA 79, 063602 (2009))



- entropy density  $s = -\frac{\partial U}{\partial T}$ ,
- energy density  $\epsilon = -p + Ts + \mu n$ ,
- specific heat  $c_v$ ,
- isoth. compressibility  $\kappa_T$ ,
- adiab. compressibility  $\kappa_S$ ,
- velocity of sound I,
- velocity of sound II,
- correlation length.

## *Fermi gases with different physics*

- 1 component Fermi gas - no s-wave interaction
- 2 component Fermi gas - BCS-BEC crossover
- 3 component Fermi gas - ??
  - Three-body problem: Efimov effect  
(Efimov, Phys. Lett. **33B**, 563 (1970),  
Review: Braaten and Hammer, Phys. Rep. **428**, 259 (2006))
  - On the lattice: Trion formation  
(Rapp, Zarand, Honerkamp, and Hofstetter, PRL **98**, 160405 (2007),  
Rapp, Hofstetter and Zarand, PRB **77**, 144520 (2008).)

## Three component Fermi gas

- For equal masses, densities etc. global SU(3) symmetry

$$\begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} \rightarrow u \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}, \quad u \in \text{SU}(3).$$

Similar to flavor symmetry in the Standard model!

- For small scattering length  $|a| \rightarrow 0$ 
  - BCS ( $a < 0$ ) or BEC ( $a > 0$ ) superfluidity at small T.
  - order parameter is conjugate triplet  $\bar{\mathbf{3}}$  under SU(3)


$$\varphi = \begin{pmatrix} \varphi_1 \\ \varphi_2 \\ \varphi_3 \end{pmatrix} \sim \begin{pmatrix} \psi_2\psi_3 \\ \psi_3\psi_1 \\ \psi_1\psi_2 \end{pmatrix}.$$

- SU(3) symmetry is broken spontaneously for  $\varphi \neq 0$ .
- What happens for large  $|a|$ ?

## Simple truncation for fermions with three components


$$\begin{aligned}\Gamma_k = & \int_x \psi^\dagger (\partial_\tau - \vec{\nabla}^2 - \mu) \psi + \varphi^\dagger (\partial_\tau - \frac{1}{2} \vec{\nabla}^2 + m_\varphi^2) \varphi \\ & + \chi^* (\partial_\tau - \frac{1}{3} \vec{\nabla}^2 + m_\chi^2) \chi \\ & + h \epsilon_{ijk} (\varphi_i^* \psi_j \psi_k + h.c.) + g (\varphi_i \psi_i^* \chi + h.c.).\end{aligned}$$

- Units are such that  $\hbar = k_B = 2M = 1$
- Wavefunction renormalization for  $\psi$ ,  $\varphi$  and  $\chi$  is implicit.
- $\Gamma_k$  contains terms for
  - fermion field  $\psi = (\psi_1, \psi_2, \psi_3)$
  - bosonic field  $\varphi = (\varphi_1, \varphi_2, \varphi_3) \sim (\psi_2 \psi_3, \psi_3 \psi_1, \psi_1 \psi_2)$
  - trion field  $\chi \sim \psi_1 \psi_2 \psi_3$




## “Refermionization”

- Trion field is introduced via a generalized Hubbard-Stratonovich transformation





- Fermion-boson coupling is regenerated by the flow



- Express this again by trion exchange  
(Gies and Wetterich, PRD **65**, 065001 (2002),  
Floerchinger and Wetterich, PLB **680**, 371 (2009).)

# Binding energies


- Vacuum limit  $T \rightarrow 0$ ,  $n \rightarrow 0$ .



- Binding energy per atom for
  - molecule or dimer  $\varphi$  (dashed line)
  - trion or trimer  $\chi$  (solid line)
- For large scattering length  $a$  trion is energetically favorable!
- Three-body bound state even for  $a < 0$ .



# Quantum phase diagram




- BCS-Trion-BEC transition

(Floerchinger, Schmidt, Moroz and Wetterich, PRA **79**, 013603 (2009)).

- $a \rightarrow 0_-$ : Cooper pairs,  $SU(3) \times U(1) \rightarrow SU(2) \times U(1)$ .
  - $a \rightarrow 0_+$ : BEC of molecules,  $SU(3) \times U(1) \rightarrow SU(2) \times U(1)$ .
  - $a \rightarrow \pm\infty$ : Trion phase,  $SU(3)$  unbroken.
- Quantum phase transitions
    - from BCS to Trion phase
    - from Trion to BEC phase.

# Efimov effect



- Self-similarity in energy spectrum.
- Efimov trimers become more and more shallow. At  $a = \infty$

$$E_{n+1} = e^{-2\pi/s_0} E_n.$$


- Simple truncation:  $s_0 \approx 0.82$ .
- Advanced truncation:  $s_0 \approx 1.006$  (exact result)  
(Moroz, Floerchinger, Schmidt and Wetterich, PRA **79**, 042705 (2009).)

## Renormalization group limit cycle

- For  $\mu = 0$  and  $a^{-1} = 0$  flow equations for rescaled couplings

$$k \frac{\partial}{\partial k} \begin{pmatrix} \tilde{g}^2 \\ \tilde{m}_\chi^2 \end{pmatrix} = \begin{pmatrix} 7/25 & -13/25 \\ 36/25 & 7/25 \end{pmatrix} \begin{pmatrix} \tilde{g}^2 \\ \tilde{m}_\chi^2 \end{pmatrix}.$$


- Solution is log-periodic in scale.



- Every zero-crossing of  $\tilde{m}_\chi^2$  corresponds to a new bound state.
- For  $\mu \neq 0$  or  $a^{-1} \neq 0$  limit cycle scaling stops at some scale  $k$ . Only finite number of Efimov trimers.

## Contact to experiments



- Model can be generalized to case without SU(3) symmetry (Floerchinger, Schmidt and Wetterich, PRA A **79**, 053633 (2009)).
- Hyperfine states of  ${}^6\text{Li}$  have large scattering lengths.



- Binding energies might be measured using RF-spectroscopy.
- Lifetime is quite short  $\sim 10\text{ns}$ .

## Three-body loss rate

- Three-body loss rate measured experimentally (Ottenstein et al., PRL **101**, 203202 (2008); Huckans et al., PRL **102**, 165302 (2009))



- Trion may decay into deeper bound molecule states
- Calculate  $B$ -field dependence of loss process above.
- Left resonance (position and width) fixes model parameters.
- Form of curve for large  $B$  is prediction.
- Similar results obtained by other methods (Braaten, Hammer, Kang and Platter, PRL **103**, 073202 (2009); Naidon and Ueda, PRL **103**, 073203 (2009).)

# Conclusions

- Functional renormalization is a useful method to describe ultracold quantum gases.
- Quantitative precision seems reachable.
- Unified description of
  - Bosons and Fermions,
  - Weak and strong coupling,
  - Few-Body and Many-Body physics.