

Threshold resummation of heavy coloured particle cross sections

Pietro Falgari

Institute for Theoretical Physics, Universiteit Utrecht

“Strong interactions: From methods to structures”
12-16 February 2011


Based on [M. Beneke, PF, S. Klein, C. Schwinn](#) [Nucl. Phys. B828: 69-101, 2010],
[Nucl. Phys. B842: 414-474, 2011] and work in preparation

Motivation


IN THIS TALK: pair production of **coloured heavy particles** at Tevatron/LHC

$$N_1 N_2 \rightarrow H(p_1) H'(p_2) + X \quad H, H' = \text{top, squarks, gluinos...}$$

accurate theoretical predictions for the cross section phenomenologically important
(sensitivity to **mass parameters, exclusion bounds, model discrimination...**)



[D0 Collaboration '09]




[Plehn, PROSPINO]

+ theoretically interesting due to **non-trivial colour exchange**

Soft-gluon and Coulomb corrections

NLO partonic cross sections enhanced near **threshold**, $\beta \equiv \sqrt{1 - (m_H + m_{H'})^2/\hat{s}} \rightarrow 0$

- **Threshold logarithms:** $\sim \alpha_s^n \ln^m \beta$
 \Leftrightarrow **soft-gluon exchange** between initial-initial, initial-final ($\alpha_s \ln^{2,1} \beta$) and final-final state particles ($\alpha_s \ln \beta$)
- **Coulomb corrections:** $\sim (\alpha_s/\beta)^n$
 \Leftrightarrow **static interaction** of slowly-moving heavy particles (mediated by **potential gluons**...)



enhanced terms can spoil convergence of perturbative series \Rightarrow **RESUMMATION**

- \Rightarrow **normalisation** of the cross section
- \Rightarrow reduction of dependence on the **factorisation-scale**
- \Rightarrow can be used to construct **higher-order approximations at fixed order in α_s**

● $t\bar{t}$ production

- **NLO QCD:** Nason et al. '88; Beenakker et al. '89
- **NLO EW:** Beenakker et al. '94; Bernreuther et al. '95; Kuhn et al. '96;...
- **NNLO:** in progress Bonciani et al. '10; Czakon '11
- **NLL (+NLO):** Kidonakis et al. '96; Bonciani et al. '98; Cacciari et al. '08; Moch et al. '08; Kidonakis et al. '08;...
- **NNLL, approx. NNLO:** Beneke, PF, Klein, Schwinn '09/'10; Ahrens et al. '10; Kidonakis '10; HATHOR Aliev et al. '10

● Squarks, gluinos

- **NLO SUSY-QCD:** Beenakker et al. '96; PROSPINO, Plehn et al.
- **NLO EW:** Bornhauser et al. '07; Hollik et al. '07-'10; Gerner et al '10
- **NLL/approx. NNLO:** Beneke, PF, Schwinn '10; Kulesza/Motyka '09; Beenakker et al. '09/'10; Langefeld/Moch '09/'10

+ many works on **Coulomb resummation** (\Leftrightarrow quarkonia physics, $e^-e^+ \rightarrow t\bar{t}, \dots$)


Combined soft and Coulomb resummation

$\alpha_s/\beta \sim \alpha_s \ln \beta \sim 1 \Rightarrow$ modified counting scheme

$$\hat{\sigma}_{pp'} \propto \hat{\sigma}^{(0)} \sum_{k=0} \left(\frac{\alpha_s}{\beta} \right)^k \exp \left[\underbrace{\ln \beta g_0(\alpha_s \ln \beta)}_{(\text{LL})} + \underbrace{g_1(\alpha_s \ln \beta)}_{(\text{NLL})} + \underbrace{\alpha_s g_2(\alpha_s \ln \beta)}_{(\text{NNLL})} + \dots \right] \\ \times \left\{ 1 (\text{LL,NLL}); \alpha_s, \beta (\text{NNLL}); \alpha_s^2, \alpha_s \beta, \beta^2 (\text{NNNLL}); \dots \right\}$$

- **non-relativistic H, H' and Coulomb gluons:**
 $E \sim m_H \beta^2, |\vec{p}| \sim m_H \beta$
- **soft gluons:** $q_s \sim m_H \beta^2$

potential and soft modes have the same energy and can “communicate” with each other




\Rightarrow **structure of soft-Coulomb emission can be in principle highly non-trivial!**

Factorisation of pair production near threshold

Effective-theory description of pair production near threshold $\hat{s} \sim (m_H + m_{H'})^2$

[Benke, PF, Schwinn, '09/'10] \Rightarrow factorization of **hard**, **soft** and **Coulomb** contributions

$$\hat{\sigma}_{pp'}(\hat{s}, \mu_f) = \sum_i H_i(M, \mu_f) \int d\omega \sum_{R_\alpha} J_{R_\alpha}(E - \frac{\omega}{2}) W_i^{R_\alpha}(\omega, \mu_f)$$



- hard function H_i depends on the **specific physics model and process**
- potential function J_{R_α} encodes Coulomb effects ($\sim \alpha_s^n / \beta^n$)
- process-independent soft function $W_i^{R_\alpha}$ ($\sim \alpha_s^n \ln^m \beta$)
 \Rightarrow depends only on total colour charge R_α of the pair!

factorization valid up to NNLL and for S-wave production

EFT description of pair-production near threshold

Near threshold ($\beta \ll 1$) partonic cross section receives contributions from four different momentum regions ($M \equiv m_H + m_{H'}$):

- **hard**: $k^2 \sim M^2$
- **(ultra)-soft**: $k_0 \sim |\vec{k}| \sim M\beta^2$
- **potential**: $k_0 \sim M\beta^2, |\vec{k}| \sim M\beta$
- **collinear**: $k_- \sim M, k_+ \sim M\beta^2, k_\perp \sim M\beta$

full theory matched on an effective Lagrangian from which **hard modes** are integrated out.

$$\mathcal{L}_{\text{full}} \rightarrow \mathcal{L}_{\text{EFT}} \equiv \mathcal{L}_{\text{SCET}} + \mathcal{L}_{\text{PNRQCD}}$$

- $\mathcal{L}_{\text{SCET}}$: describes interactions of **collinear** (ξ_c, A_c) and **soft** (A_s) modes

$$\mathcal{L}_{\text{SCET}} = \bar{\xi}_c \left(in \cdot D + i\not{D}_\perp c \frac{1}{i\bar{m} \cdot D_c} i\not{D}_\perp c \right) \frac{\not{n}}{2} \xi_c - \frac{1}{2} \text{tr} \left(F_c^{\mu\nu} F_{c\mu\nu} \right) + \dots$$

- $\mathcal{L}_{\text{PNRQCD}}$: contains interactions of **potential** (ψ, ψ') and **soft** (A_s) modes

$$\begin{aligned} \mathcal{L}_{\text{PNRQCD}} = & \psi^\dagger \left(iD_s^0 + \frac{\vec{\partial}^2}{2m_H} + \frac{i\Gamma_H}{2} \right) \psi + \psi'^\dagger \left(iD_s^0 + \frac{\vec{\partial}^2}{2m_{H'}} + \frac{i\Gamma_{H'}}{2} \right) \psi' \\ & + \int d^3\vec{r} \left[\psi^\dagger \mathbf{T}^{(R)a} \psi \right] (x + \vec{r}) \left(\frac{\alpha_s}{r} \right) \left[\psi'^\dagger \mathbf{T}^{(R')a} \psi' \right] (x) + \dots \end{aligned}$$

Structure of EFT amplitudes

$$\mathcal{A}(pp' \rightarrow HH'X) = \sum_{\ell} C_{\{a;\alpha\}}^{(\ell)}(\mu_f) \langle HH'X | \mathcal{O}_{\{a;\alpha\}}^{(\ell)}(\mu_f) | pp' \rangle_{\text{EFT}}$$

- effective operators $\mathcal{O}_{\{a;\alpha\}}^{(0)}(\mu_f) \propto [\phi_{c;a_1,\alpha_1} \phi_{\bar{c};a_2,\alpha_2} \psi_{a_3\alpha_3}^{\dagger} \psi_{a_4\alpha_4}^{\dagger}]$ contain collinear and non-relativistic fields \Leftrightarrow **long-distance** effects.
Operators with more fields or derivatives suppressed by extra powers of β (not required at NNLL...)
- matrix element evaluated using the EFT Lagrangian \Rightarrow soft gluons interacting with everything and potential interactions between the two non-relativistic heavy particles
- hard matching coefficient $C_{\{a;\alpha\}}^{(\ell)}(\mu_f)$ encodes **short-distance** structure of pair-production process at the scale M
 - \Rightarrow extracted from fixed-order calculations of on-shell amplitudes
 - \Rightarrow decomposed on a suitable basis of colour-state operators:
$$C_{\{a;\alpha\}}^{(\ell)}(\mu_f) = C_{\{\alpha\}}^{(\ell,i)}(\mu_f) c_{\{a\}}^{(i)}$$

Soft-gluon decoupling

At **leading order in β** soft gluons can be decoupled from the effective Lagrangian via field redefinitions involving **soft Wilson lines (path-order exponentials of soft gluon fields)**:

$$\phi_c(x) \rightarrow S_n^{(R)}(x_-)\phi_c^{(0)}(x)$$

$$\psi(x) \rightarrow S_v^{(R)}(x_0)\psi^{(0)}(x) \quad S_n^{(R)}(x) = \text{P exp} \left[ig_s \int_{-\infty}^0 dt n \cdot A_s^c(x + nt) \mathbf{T}^{(R)c} \right]$$

$$S_v^{(R)\dagger}(x_0) D_s^0 S_v^{(R)}(x_0) = \partial^0 \quad [\psi^\dagger \mathbf{T}^{(R)a} \psi](x + \vec{r}) = S_{v,ab}^8(x_0) [\psi^{(0)\dagger} \mathbf{T}^{(R)b} \psi^{(0)}](x + \vec{r})$$

upon field redefinition:

$$\hat{\sigma}_{pp'}(\hat{s}, \mu_f) \equiv \frac{1}{2s} \int d\Phi |\mathcal{A}|^2 = \sum_{i,i'} \sum_{S=|s-s'|}^{s+s'} \underbrace{H_{ii'}^S(M, \mu_f)}_{\text{hard}} \int d\omega \sum_{R_\alpha} \underbrace{J_{R_\alpha}^S(E - \frac{\omega}{2})}_{\text{potential}} \underbrace{W_{ii'}^{R\alpha}(\omega, \mu_f)}_{\text{soft}}$$

- $H_{ii'}^S(M, \mu_f) \propto C_{\{\alpha\}}^{(0,i)}(M, \mu_f) C_{\{\beta\}}^{(0,i')*}(M, \mu_f) \dots$
- $J_{R_\alpha}^S(q) \propto \int d^4z e^{iq \cdot z} \langle 0 | [\psi'^{(0)} \psi^{(0)}](z) [\psi^{(0)\dagger} \psi'^{(0)\dagger}](0) | 0 \rangle$
- $W_{ii'}^{R\alpha}(\omega, \mu_f) = P_{\{k\}}^{R\alpha} c_{\{a\}}^{(i)} c_{\{b\}}^{(i')} \int dz_0 e^{i\omega z_0/2} \langle 0 | \bar{\mathbf{T}}[S_n^\dagger S_n^\dagger S_v S_v](z) \mathbf{T}[S_{\bar{n}} S_n S_v^\dagger S_v^\dagger](0) | 0 \rangle$

Colour structure of the factorisation formula

The factorisation formula has a priori a **non-trivial colour structure**

- hard function is a matrix in colour-state space: $H_{ii'} \equiv H_{\{ab\}} c_{\{a\}}^{(i)} c_{\{b\}}^{(i')*}$
- potential function $J_{\{k\}}$ is projected over irreducible representations of the HH' system:
 $J_{\{k\}} = \sum_{R_\alpha} P_{\{k\}}^{R_\alpha} J_{R_\alpha}$, with $R \otimes R' = \sum_{\alpha} R_\alpha$
- soft function given by a set of colour matrices $W_{ii'}^{R_\alpha}$

Colour basis $c_{\{a\}}^{(i)}$ can be chosen such that $W_{ii'}^{R_\alpha}$ are diagonal to all orders in α_s

[Beneke, PF, Schwinn, Nucl.Phys. B828 (2010)]

- ⇒ decompose initial-state and final-state product representations into **irreducible representations**:
→ Clebsch-Gordan coefficients

$$r \otimes r' = \sum_{\alpha} r_{\alpha} \rightarrow C_{\alpha a_1 a_2}^{r_{\alpha}} \qquad R \otimes R' = \sum_{\beta} R_{\beta} \rightarrow C_{\alpha a_1 a_2}^{R_{\beta}}$$

- ⇒ identify pairs of equivalent initial- and final-state representations $P_i = (r_{\alpha}, R_{\beta})$
⇒ construct colour basis by contracting the Clebsches into colour-invariant combinations

$$c_{\{a\}}^{(i)} = \frac{1}{\sqrt{\dim(r_{\alpha})}} C_{\alpha a_1 a_2}^{r_{\alpha}} C_{\alpha a_3 a_4}^{R_{\beta}*} \qquad P_{\{a\}}^{R_{\alpha}} = C_{\alpha a_1 a_2}^{R_{\alpha}*} C_{\alpha a_3 a_4}^{R_{\alpha}}$$

Soft/hard resummation in momentum space


IR structure of QCD amplitudes and scale-invariance of the hadronic cross section give RG evolution equations for the soft function $W_i^{R\alpha}$ and the hard function $H_i^{R\alpha}$
(generalisation of DY result [Becher, Neubert, Xu '07] to arbitrary R_α)

$$\frac{d}{d \ln \mu_f} W_i^{R\alpha}(\omega, \mu_f) = -2 \left[(C_r + C_{r'}) \Gamma_{\text{cusp}} \ln \left(\frac{\omega}{\mu_f} \right) + 2\gamma_{H,S}^{R\alpha} + 2\gamma_s^r + 2\gamma_s^{r'} \right] W_i^{R\alpha}(\omega, \mu_f) \\ - 2(C_r + C_{r'}) \Gamma_{\text{cusp}} \int_0^\omega d\omega' \frac{W_i^{R\alpha}(\omega', \mu_f) - W_i^{R\alpha}(\omega, \mu_f)}{\omega - \omega'}$$

and similar for hard function $H_i(M, \mu_f)$

Resummation strategy


- Solve evolution equation in momentum space
- Evolve the function H_i from the hard scale μ_h to μ_f
- Evolve soft function $W_i^{R\alpha}$ from a low scale μ_s to μ_f .



Resummation of Coulomb corrections

Exchange of Coulomb gluons between the pair H, H' : $\Delta\sigma^{\text{Coul},(1)}/\sigma^{\text{tree}} \sim \alpha_s/\beta \sim 1$

\Rightarrow **Coulomb corrections must be resummed to all orders as well**



Resummation of Coulomb effects well understood from **PNRQCD** and quarkonia physics.

For HH' system in **irreducible representation R_α** (and at LO in PNRQCD):

$$J_{R_\alpha}(E) = -\frac{(2m_{\text{red}})^2}{2\pi} \text{Im} \left\{ \sqrt{-\frac{E}{2m_{\text{red}}}} + \alpha_s(-D_{R_\alpha}) \left[\frac{1}{2} \ln \left(-\frac{8m_{\text{red}}E}{\mu_f^2} \right) - \frac{1}{2} + \gamma_E + \psi \left(1 - \frac{\alpha_s(-D_{R_\alpha})}{2\sqrt{-E/(2m_{\text{red}})}} \right) \right] \right\} \quad E \equiv \sqrt{s} - M$$

+ higher-order Coulomb and non-Coulomb potentials at NNLL!

Squark-antisquark production at the LHC

$$PP \rightarrow \tilde{q}\tilde{q} + X$$

NLL soft resummation and Coulomb resummation to total cross section

$$\hat{\sigma}_{pp'}^{\text{NLL}}(\hat{s}, \mu_f) = \sum_i H_i(\mu_h) U_i(M, \mu_h, \mu_s, \mu_f) \frac{e^{-2\gamma_E\eta}}{\Gamma(2\eta)} \int_0^\infty d\omega \frac{J_{R_\alpha}(E - \frac{\omega}{2})}{\omega} \left(\frac{\omega}{2M}\right)^{2\eta}$$

resummed cross section is matched onto the full NLO result

[Zerwas et al., '96; Langenfeld, Moch '09]

$$\hat{\sigma}_{pp'}^{\text{match}}(\hat{s}, \mu_f) = \left[\hat{\sigma}_{pp'}^{\text{NLL}}(\hat{s}, \mu_f) - \hat{\sigma}_{pp'}^{\text{NLL}}(\hat{s}, \mu_f)|_{\text{NLO}} \right] + \hat{\sigma}_{pp'}^{\text{NLO}}(\hat{s}, \mu_f)$$

Scale choice for μ_s , μ_h and μ_C

What is a good choice for μ_s , μ_h and μ_C ?

- **Hard scale:** $\mu_h = 2m_{\bar{q}}$
- Choose **soft scale** such that one-loop soft corrections to the **hadronic cross section** are minimised [Becher, Neubert, Xu '07]

$$\frac{\partial}{\partial \bar{\mu}_s} \int dx_1 dx_2 f(x_1, \bar{\mu}_s) f(x_2, \bar{\mu}_s) \Delta \hat{\sigma}^{S,(1)}(\hat{s}, \bar{\mu}_s) = 0$$

This choice guarantees well-behaved perturbative expansion at the low scale $\bar{\mu}_s$

- **Coulomb scale:** set by typical **virtuality of a Coulomb gluons** $\sqrt{|q^2|} \sim m_{\bar{q}}\beta \sim m_{\bar{q}}\alpha_s$

$$\Rightarrow \mu_C = \max\{2m_{\bar{q}}\beta, C_F m_{\bar{q}}\alpha_s(\mu_C)\}$$

\hookrightarrow twice **inverse Bohr radius** of first bound state

Necessary to correctly resum NLL effects from running of Coulomb potential!


Squark-antisquark resummed cross section

Beneke, PF, Schwinn '10

- **NLL**: full soft and Coulomb resummation (including bound-state contributions from below threshold)
- **NLL_{noBS}**: soft and Coulomb resummation (but no bound-state contribution)
- **NLL_{s+h} + C**: soft resummation + Coulomb resummation (no interference terms)
- **NLL_{s+h}**: soft resummation only


Setup:

- PP@ 14 TeV
- MSTW2008 PDFs
- equal squark masses
- no stops
- $m_{\tilde{g}} = 1.25m_{\tilde{q}}$
- $\mu_f = m_{\tilde{q}}$




Factorisation-scale dependence

One of the motivations for resummation is **reduction of scale dependence** of NLO result:



LHC@14TeV



$m_{\tilde{q}} = 1 \text{ TeV}$

All scales varied by a factor 2 around the default values, and uncertainties summed in quadrature

Comparison with Mellin-space resummation

[**Beenakker et al. JHEP 0912:041, 2009**] (**Mellin-space formalism**, only soft resummation, no Coulomb effects) \Rightarrow **compare to our approximation NLL_{s+h} with $\mu_h = \mu_f$**

$m_{\tilde{q}}$ [GeV]	$\sigma(pp \rightarrow \tilde{q}\tilde{q})(\text{pb}), \sqrt{s} = 14 \text{ TeV}$			
	NLO	NLL_{Mellin}	NLL_s	NLL
200	1.3×10^3	1.31×10^3 (1%)	1.31×10^3 (1%)	1.34×10^3 (3.4%)
500	1.6×10^1	1.61×10^1 (1.2%)	1.62×10^1 (1.3%)	1.67×10^1 (4.2%)
1000	2.89×10^{-1}	2.93×10^{-1} (1.7%)	2.94×10^{-1} (1.7%)	3.06×10^{-1} (5.8%)
2000	1.11×10^{-3}	1.14×10^{-3} (3.4%)	1.14×10^{-3} (3.1%)	1.24×10^{-3} (11%)
3000	7.13×10^{-6}	7.59×10^{-6} (6.4%)	7.54×10^{-6} (5.8%)	8.61×10^{-6} (21%)

- Good agreement of momentum-space and Mellin-moment resummation
- **Full soft-Coulomb resummation generally larger than pure soft resummation!**

$t\bar{t}$ production at NNLL/NNLO


All ingredients for **NNLL resummation** of $t\bar{t}$ cross section known

- 1-loop colour-separated hard functions $H_i^{(1)}$ [Czakon, Mitov '09]
- **2-loop soft anomalous dimension** [Beneke, PF, Schwinn '09; Czakon, Mitov, Sterman '09]
- NLO Coulomb and non-Coulomb potentials [Beneke, Signer, Smirnov '99]

Can be used to construct approx. NNLO containing all terms singular in β

[Beneke, PF, Czakon, Mitov, Schwinn '09; HATHOR Aliev et al. '10]

$$\begin{aligned}\hat{\sigma}_{\text{approx.}}^{\text{NLO}} &= \frac{k_{\text{LO}}^2}{\beta^2} + \frac{1}{\beta} [k_{\text{NLO},1} \ln \beta + k_{\text{NLO},0}] + k_{\text{n-C}} \ln \beta \\ &+ c_{S,4}^{(2)} \ln^4 \beta + c_{S,3}^{(2)} \ln^3 \beta + c_{S,2}^{(2)} \ln^2 \beta + c_{S,1}^{(2)} \ln \beta \\ &+ H^{(1)} [c_{S,2}^{(1)} \ln^2 \beta + c_{S,1}^{(1)} \ln \beta] \\ &+ \frac{k_{\text{LO}}}{\beta} [c_{S,2}^{(1)} \ln^2 \beta + c_{S,1}^{(1)} \ln \beta + c_{S,0}^{(1)} + H^{(1)}]\end{aligned}$$



NNLL/NNLO total $t\bar{t}$ cross section

$m_t = 173.1$ GeV, $\mu_f = m_t$, MSTW2008NNLO

Beneke, PF, Klein, Schwinn, **PRELIMINARY!**

$\sigma_{t\bar{t}}$ [pb]	Tevatron	LHC@7	LHC@10	LHC@14
NLO	6.50 ^{+0.32+0.33} _{-0.70-0.24}	150 ⁺¹⁸⁺⁸ ₋₁₉₋₈	380 ⁺⁴⁴⁺¹⁷ ₋₄₆₋₁₇	842 ⁺⁹⁷⁺³⁰ ₋₉₇₋₃₂
NLO+NLL	6.57 ^{+0.52+0.33} _{-0.30-0.24}	151 ⁺²³⁺⁸ ₋₁₂₋₈	382 ⁺⁶⁰⁺¹⁷ ₋₃₂₋₁₈	848 ⁺¹³⁶⁺³⁰ ₋₇₅₋₃₂
NLO+NNLL	6.77 ^{+0.27+0.35} _{-0.48-0.25}	155 ⁺⁴⁺⁸ ₋₉₋₉	390 ⁺¹⁴⁺¹⁷ ₋₂₆₋₁₈	858 ⁺³⁵⁺³¹ ₋₆₄₋₃₃
NNLO _{app} (β)	7.10 ^{+0.0+0.36} _{-0.26-0.26}	162 ⁺²⁺⁹ ₋₃₋₉	407 ⁺⁹⁺¹⁷ ₋₅₋₁₈	895 ⁺²⁴⁺³¹ ₋₆₋₃₃
NNLO_{app}(β)+NNLL	7.13 ^{+0.22+0.36} _{-0.24-0.26}	162 ⁺⁴⁺⁹ ₋₁₋₉	405 ⁺¹⁴⁺¹⁷ ₋₂₋₁₈	892 ⁺³⁸⁺³¹ ₋₃₋₃₃
NNLO_{app}(β)+NNLL+BS	7.14 ^{+0.14+0.36} _{-0.22-0.26}	162 ⁺⁴⁺⁹ ₋₁₋₉	407 ⁺¹⁴⁺¹⁷ ₋₂₋₁₈	896 ⁺³⁸⁺³¹ ₋₃₋₃₃

- Combined soft-Coulomb resummation for $t\bar{t}$ total cross section
- All scales ($\mu_f, \mu_h, \mu_s, \mu_C$) varied in interval $0.5\tilde{\mu}_i < \mu_i < 2\tilde{\mu}_i$
- Fixed μ_s from minimising $\Delta\sigma_{\text{soft}}^{\text{NLO}}$: $\Rightarrow \mu_s = 85, 146, 174, 202$ GeV for Tevatron, LHC@7, LHC@10, LHC@14. **No large scale hierarchy**
- Residual scale uncertainty for NNLO_{app}(β)+NNLL+BS $\sim 5\%$
+ **estimated $\sim 3\%$ uncertainty from $\alpha_s^2 H_i^{(2)}$ contribution**

- Factorisation formula for pair-production near threshold in SCET+PNRQCD
 - ⇒ Valid for **arbitrary colour representations**
 - ⇒ Proves decoupling of **hard, soft** and **Coulomb** modes
 - ⇒ **Diagonal in colour-space** to all orders in α_s
- Simultaneous resummation of threshold logarithms and Coulomb singularities
 - ⇒ Directly in **momentum space** via RG evolution equations
- Application to squark-antisquark production at the LHC
 - ⇒ NLL corrections $\sim 4 - 20\%$ for $m_{\tilde{q}} \sim 200\text{GeV} - 3\text{TeV}$
 - ⇒ Reduction of factorisation-scale dependence
- NNLL resummation of $t\bar{t}$ total cross section
 - ⇒ All $O(\alpha_s^2)$ **terms singular in β** included
 - ⇒ NNLL corrections beyond NNLO very small

Backup slides

HO Coulomb and non-Coulomb corrections in PNRQCD required at NNLL/NNLO

- **Coulomb potential:**

$$\tilde{V}_C^{(1)}(\vec{p}, \vec{q}) = \frac{D_{R_\alpha} \alpha_s^2}{|\vec{q}|^2} \left(a_1 - \beta_0 \ln \frac{|\vec{q}|}{\mu^2} \right)$$

- **Non-Coulomb potentials:**

$$\tilde{V}_{nC}^{(1)}(\vec{p}, \vec{q}) = \frac{4\pi D_{R_\alpha} \alpha_s}{|\vec{q}|^2} \left[\frac{\pi \alpha_s |\vec{q}|}{4m} \left(\frac{D_{R_\alpha}}{2} + C_A \right) + \frac{|\vec{p}|^2}{m^2} + \frac{|\vec{q}|^2}{m^2} v_{\text{spin}} \right]$$

$$v_{\text{spin}} = 0(\text{singlet}), -\frac{2}{3}(\text{triplet})$$

contribution to NNLO total cross section

$$\Delta\sigma_{nC}^{\text{NNLO}} = \sigma^{(0)} \alpha_s^2 \ln \beta \left[D_{R_\alpha}^2 (1 + v_{\text{spin}}) + D_{R_\alpha} \right] C_A$$


Subleading soft interactions

At NNLL subleading soft vertices in SCET and PNRQCD potentially important

$$\psi^\dagger \vec{x} \cdot \vec{E}_{\text{us}}(x_0, \vec{0}) \psi \quad \bar{\xi} \left(x_\perp^\mu n_-^\nu W_c g F_{\mu\nu}^{us} W_c^\dagger \right) \frac{\not{n}_+}{2} \xi$$

Subleading soft interactions not removed by field redefinitions

⇒ related to off-diagonal three-parton colour correlations in IR singularities of QCD amplitudes (Ferroglia, Neubert, Pecjak, Yang '09)



$$\Rightarrow \frac{\alpha_s}{\beta} \alpha_s \beta \ln \beta \sim \alpha_s^2 \ln \beta$$

Contributions of subleading soft-collinear and soft-potential vertices vanish for the total cross section!


- **Soft-collinear:** k_\perp can always be chosen to be 0
- **Soft-potential:** vanish because of rotational invariance

Contribution of threshold-enhanced terms

At LHC $\sqrt{s} \gg 2m_t \Rightarrow$ **How good is the threshold approximation?**
can study the approximation at the NLO level...

Plot $8\beta m_t^2 / (s(1 - \beta^2)^2) \mathcal{L}_{gg}(\beta) \hat{\sigma}_{tt}(\beta)$:

- **NLO:** exact NLO result
- **NLO sing.:** only singular terms in β
- **NLO approx.:** singular terms + $O(1)$ term ($\Leftrightarrow H_i^{(1)}$)



NLO sing. is good approximation only up to $\beta \sim 0.3$

However: expect NNLO approximation to be better (more singular terms at $O(\alpha_s^2)$)

Alternative approaches

- **Pair invariant-mass distribution** $d\sigma(\bar{t}\bar{t} + X)/dM_{\bar{t}\bar{t}}$
 [Kidonakis, Sterman '97; Ahrens et al. '10]

$$\left[\frac{\ln^n(1-z)}{(1-z)} \right]_+ \quad z = \frac{M_{\bar{t}\bar{t}}^2}{\hat{s}}$$

- **One-particle inclusive cross section** $d\sigma(t + X)/ds_4$:
 [Laenen, Oderda, Sterman '98; Kidonakis '10]

$$\left[\frac{\ln^n(s_4/m_t^2)}{s_4} \right] \quad s_4 = p_X^2 - m_t^2$$

	$\sigma_{\bar{t}\bar{t}}[\text{pb}]$	Tevatron	LHC@7	LHC@10	LHC@14
BFKS	NLO	$6.50^{+0.32+0.33}_{-0.70-0.24}$	150^{+18+8}_{-19-8}	380^{+44+17}_{-46-17}	842^{+97+30}_{-97-32}
	NLO+NNLL	$6.77^{+0.27+0.35}_{-0.48-0.25}$	155^{+4+8}_{-9-9}	390^{+14+17}_{-26-18}	858^{+35+31}_{-64-33}
	NNLO(β)	$7.10^{+0.0+0.36}_{-0.26-0.26}$	162^{+2+9}_{-3-9}	407^{+9+17}_{-5-18}	895^{+24+31}_{-6-33}
Kidonakis '10	NNLO(β)	$7.08^{+0.0+0.36}_{-0.24-0.27}$	163^{+7+9}_{-5-9}	415^{+17+18}_{-21-19}	920^{+50+33}_{-39-35}
Ahrens et al. '10	NLO+NNLL	$6.48^{+0.17+0.32}_{-0.21-0.25}$	146^{+7+8}_{-7-8}	368^{+20+19}_{-14-15}	813^{+50+30}_{-36-35}
	NNLO(β)	$6.55^{+0.32+0.33}_{-0.41-0.24}$	149^{+10+8}_{-9-8}	377^{+28+16}_{-23-18}	832^{+65+31}_{-50-29}